arXiv:cond-mat/9808167v1 [cond-mat.str-el] 16 Aug 1998

SO(N) symmetries in the two–chain model of correlated fermions

H.J. Schulz
Laboratoire de Physique des Solides, Université Paris–Sud
91405 Orsay,France

The two–chain model of correlated fermions, appropriate e.g. for doped spin ladders or carbon nanotubes, is investigated in the weak–interaction limit. It is shown that the model scales to situations with high internal symmetries, generically SO(6), whereas on critical lines there is SO(5) or SO(3) symmetry. Consequences for the level spectroscopy of the model are discussed. In particular, the "π–mode" is shown to be sixfold degenerate and contains both spin and charge excitations.

Models of two coupled chains of interacting fermions are of interest for a variety of reasons: they are directly applicable to interesting experimental systems like doped spin ladders [1] or carbon nanotubes [2], and they also provide a first step for the understanding of the transition from the well–understood single chain case to the area of two–dimensional correlated fermions. [3] In particular, both analytical [4–6] and numerical [7–9] work show that the two–chain system with purely repulsive interactions is a d–type superconductor whereas the single chain is antiferromagnetic. This provides a potential explanation for the observation of superconductivity in a doped spin ladder, [1] and is probably the best–established case of correlation–induced superconductivity.

In recent work [10] it has been shown that the Mott–insulating half–filled two–chain model has a very big SO(8) symmetry in the weak–correlation limit, leading to a number of interesting predictions. Away from half–filling the SO(8) symmetry is broken and the system becomes conducting. Here I will study the symmetries of the conducting state away from half–filling and show that generically a still large SO(6) (=SU(4)) symmetry remains. On critical lines the symmetry is lower, SO(3) or SO(5). There are exact solutions corresponding to these symmetric models, [11,12] and from those it is possible to obtain quite detailed information about the excitation spectrum and in particular the appearance and location of bound states. The present work also provides a microscopic extension of the SO(5) model of correlated fermions [13] and the associated ladder models. [14]

I consider the standard two–chain model of interacting fermions. [4–6] After linearizing the electron dispersion around the Fermi energy, the kinetic energy density takes the form ,

$$h_0 = -iv_F \sum_{s,q=0,\pi} \{\psi^\dagger_{Rsq}\partial_x\psi_{Rsq} - \psi^\dagger_{Lsq}\partial_x\psi_{Lsq}\} \quad (1)$$

Here $q = 0$ (π) for the (anti–)bonding bands split by the interchain hopping integral $t_\perp$, s is the spin, R (L) refers to the right– (left–)going fermions, and $\psi_{R(L)sq}$ are the corresponding fermion field operators.

Important interaction effects arise from processes where both the two incoming and the two outgoing particles are all at the Fermi energy. The Hamiltonian density of the interaction then is

$$\begin{aligned} h_{int} &= g_{1,abcd}\psi^\dagger_{Rsa}\psi^\dagger_{Ltb}\psi_{Rtc}\psi_{Lsd} \\ &+ g_{2,abcd}\psi^\dagger_{Rsa}\psi^\dagger_{Ltb}\psi_{Ltc}\psi_{Rsd} \end{aligned} \quad (2)$$

These terms describe backward (g_1) and forward (g_2) scattering between particles at the Fermi surface. Summation over the spin indices s, t and the band indices a, b, c, d is understood. The interactions allowed by momentum and energy conservation are $g_{1,0000} = g_{1,\pi\pi\pi\pi} \equiv g_{11}$, $g_{1,00\pi\pi} = g_{1,\pi\pi00} \equiv g_{12}$, $g_{1,0\pi0\pi} = g_{1,\pi0\pi0} \equiv g_{13}$, $g_{2,0000} = g_{2,\pi\pi\pi\pi} \equiv g_{21}$, $g_{2,0\pi\pi0} = g_{2,\pi00\pi} \equiv g_{22}$, $g_{2,00\pi\pi} = g_{2,\pi\pi00} \equiv g_{23}$.

The one–loop renormalization group equations for the coupling constants under a change of energy cutoff $\Lambda \to \Lambda e^{-l}$ have been found previously [15] and are given by:

$$\begin{aligned} g'_{11} &= -2g^2_{11} - 2g_{12}g_{23} \\ g'_{12} &= -2(g_{12}g_{21} + g_{11}g_{23} + 2g_{12}g_{13} - g_{13}g_{23} - g_{12}g_{22}) \\ g'_{13} &= -2g^2_{12} + 2g_{12}g_{23} - 2g^2_{13} \\ g'_{21} &= -g^2_{11} - g^2_{12} - g^2_{23} \\ g'_{22} &= -g^2_{13} + g^2_{23} \\ g'_{23} &= -2g_{11}g_{12} - 2(g_{21} - g_{22})g_{23} \end{aligned} \quad (3)$$

Here $g'_{ij} \equiv 2\pi v_F \frac{dg_{ij}}{dl}$

Abelian bosonization is used to write the model in terms of symmetric and antisymmetric charge ($\nu = \rho\pm$) and spin ($\nu = \sigma\pm$) fields ϕ_ν and their conjugate momentum densities Π_ν. The total Hamiltonian becomes the sum of a term bilinear in boson fields (H_b) and a term containing nonlinear interactions $H_{\rm nl}$, with $H_b = H_{\rho+} + H_{\rho-} + H_{\sigma+} + H_{\sigma-}$,

$$H_\nu = \frac{\pi v_F}{2}\int dx \left\{(1+M_\nu)\,\Pi^2_\nu + \frac{1}{\pi^2}\,(1-M_\nu)\,(\partial_x\phi_\nu)^2\right\},$$

and $M_{\sigma\pm} = (g_{11} \pm g_{13})/2\pi v_F$,$M_{\rho\pm} = M_{\sigma\pm} - (g_{21} \pm g_{22})/\pi v_F$. The interaction term is

$$H_{\rm nl} = \frac{1}{(\pi\alpha)^2}\int dx\{(g_{23} - g_{12})\cos 2\theta_{\rho-}\cos 2\theta_{\sigma-}$$



FIG. 1. "Phase diagram" in the $g_1 - g_2$ plane ($g_{1i} \equiv g_1, g_{2i} \equiv g_2$), indicating the dominant fluctuation. CDW: charge–density wave, SCd: d–wave superconductor, OAF: orbital antiferromagnet, SCs: s–wave superconductor. On the critical lines $g_2 = 0$ and $g_1 = 2g_2$ additional critical fluctuations occur.

$$+ \; g_{23} \cos 2\theta_{\rho-} \cos 2\phi_{\sigma-} + g_{11} \cos 2\phi_{\sigma+} \cos 2\phi_{\sigma-}$$
$$+ \; g_{12} \cos 2\theta_{\rho-} \cos 2\phi_{\sigma+} - g_{13} \cos 2\phi_{\sigma+} \cos 2\theta_{\sigma-}, \quad (4)$$

where θ_ν is the field dual to ϕ_ν: $\Pi_\nu = \partial_x \theta_\nu / \pi$.

It is clearly interesting to understand the energetics and quantum numbers of the low–lying excited states of the model, in particular in order to understand the relation to numerical or experimental results. I start by investigating the generic regions of the phase diagram, fig.1, i.e. away from the critical lines $g_1 = 0$, $g_1 - 2g_2$. In particular, in the SCd region, important for Hubbard–like bare interactions, a numerical investigation shows that the asymptotic behavior near the divergence ($l \to l_c$) is given by $\tilde{g}_{ij}(l) = \tilde{g}^0_{ij}/(l_c - l)$, with the values

$$- \tilde{g}^0_{11} = \tilde{g}^0_{12} = \tilde{g}^0_{23} = 1/4, \quad \tilde{g}^0_{13} = 0,$$
$$\tilde{g}^0_{21} = -3/16, \quad \tilde{g}^0_{22} = 1/16 \; . \quad (5)$$

One can note that eq.(5) is an exact solution of the renormalization group equations (3) for a special set of initial conditions. For more general interactions, in particular the Hubbard model, the solution of the renormalization group equations tends asymptotically for $l \to l_c$ towards this solution, i.e. eq.(5) is the "fixed ray" towards which all solutions of the renormalization group equations scale in the SCd part of the phase diagram.

Interestingly, for the parameter values eq.(5) the model exhibits a high but somewhat hidden symmetry: after the particle–hole transformation on the left–going branch $\psi^\dagger_{Lsq} \leftrightarrow \psi_{L-sq}$ takes the $U(4)$ invariant form

$$H = H_0 - g \int dx (\frac{1}{4} \rho_R \rho_L + \psi^\dagger_{Rsq} \psi_{Lsq} \psi^\dagger_{Ltp} \psi_{Rtp}), \quad (6)$$

where $\rho_\alpha = \psi^\dagger_{\alpha sq} \psi_{\alpha sq}$. This is the *chiral* Gross–Neveu model, for which an exact Bethe ansatz solution exists. [12]

Another transformation, which will be more generally useful is obtained by refermionizing the bosonized Hamiltonian, eq.(4), by introducing three fermion fields $\psi_{1,2,3}$ corresponding to the $(\rho-)$, $(\sigma+)$, and $(\sigma-)$ boson fields. After a duality transformation $\phi_{\rho-} \leftrightarrow \theta_{\rho-}$ and writing the Dirac fermions in terms of hermitian (Majorana) fields χ as $\psi_{\alpha i} = \chi_{\alpha,2i-1} + i\chi_{\alpha,2i}$ the Hamiltonian in this subspace then takes the form

$$H = H_0 + 2g \int dx \left(\sum_{i=1}^{N=6} \chi_{Ri} \chi_{Li} \right)^2 , \quad (7)$$

This is the well–known Hamiltonian of the Gross–Neveu model [16] which has a SO(6) symmetry. Due to the equality of the Lie algebras so(6) and su(4), the equivalence of the chiral SU(4) and the (non–chiral) SO(6) models may seem quite natural. One should however notice that the nonchiral model has a double–degenerate symmetry–broken ground state, [16,17] whereas there is no symmetry breaking in the chiral model. This will lead to interesting consequences for the quantum numbers of the excited states. It should be emphasized that eq.(7) describes the massive part of the spectrum and that the total charge mode $\rho+$ remains massless in all cases and therefore contributes to correlation functions in an important way.

We can now use known results for the excitation spectrum of the SO(N) models [11,12] to obtain results for the excitations of the two–chain model. For the generic SU(4) case the elementary excitations are single fermions and holes with mass gap $m_0 \propto e^{-1/g}$. Due to the particle–hole transformation leading to eq.(6), right–going fermions transform as the fundamental **4** representation, whereas left–going fermions transform as the conjugate $\bar{\mathbf{4}}$. In the SO(6) language these fermionic excitations are kinks separating the two degenerate ground states, [17] whereas the SO(6) fermions correspond to two–particle excitations in the original model.

From the single–fermion gap m_0 one expects a gap $2m_0$ in two–particle excitations. This is in fact correct for all excitations created by operators of the form $\psi^\dagger_{Rsq} \psi^\dagger_{Ltq'}$. Consequently, all pairing correlation functions, irrespective of their spin (singlet or triplet) or orbital structure, decay exponentially, with *the same correlation length* $\xi = v_F/2m_0$. The one important exception of course is the "d–wave" singlet channel, which has purely algebraic decay. In the group theoretically language this corresponds to the decomposition $\mathbf{4} \otimes \bar{\mathbf{4}} = \mathbf{1} \oplus \mathbf{15}$, the **1** being the d–wave superconductor.

More structure arises in the particle–hole channel, created by operators of the type $\psi^\dagger_{Rsq} \psi_{Ltq'}$. Here the relevant decomposition is $\mathbf{4} \otimes \mathbf{4} = \mathbf{6} \oplus \mathbf{10}$, the **6** being the antisymmetric rank–2 tensor. Remarkably, the exact solution shows that particle–hole excitations in this channel are bound and appear at energy $\sqrt{2}m_0$, rather than $2m_0$. The operators creating these bound states are the spin

triplet $\psi^\dagger_{Rs0}(\sigma_\alpha)_{st}\psi_{Lt\pi}+(0\leftrightarrow\pi)$ which represents an $2k_F$ spin density wave which is out of phase between the two chains, and the "orbital triplet" $\psi^\dagger_{Rsq}(\sigma_\alpha)_{qq'}\psi_{Lsq'}$ which contains a charge density wave which is in phase between the two chains and the orbital antiferromagnet (OAF, or staggered flux phase). From the $SU(4)$ symmetry we can conclude that the gaps in these six channels are identical. It is in particular remarkable that the lowest spin and charge gap appear, due to bound state formation, at only $\sqrt{2}$ times the single–particle gap. One should notice that contrary to the triply–degenerate antiferromagnetic π–mode of the SO(5) model of correlated fermions [13], we here have an *extended π–mode with sixfold degeneracy* and which also contains charge excitations. The remaining particle–hole excitations (the **10**) all have excitation energy $2m_0$.

I now turn to the critical lines in fig.1. Both are of Ising type. For $g_1 = 0$ the asymptotics of the flow is now given by $\tilde{g}^0_{1i} = 0, \tilde{g}^0_{21} = -\tilde{g}^0_{22} = \tilde{g}^0_{23}/2 = 1/4$. One can now follow the same logic as before. However, here the self–dual combination $\cos 2\phi_{\sigma-} + \cos 2\theta_{\sigma-} \rightarrow \chi_{R3}\chi_{L3}$ appears, and consequently $\chi_{\alpha 4}$ decouples and remains massless, together with the two boson fields $\phi_{\nu+}$. The fermionic model for the massive degrees of freedom then is given by eq.(7), however now with $N = 3$. In this case the model has an $\mathrm{SU}(2)_R \times \mathrm{SU}(2)_L \times \mathrm{SU}(2)$ symmetry, where the first two subgroups represent spin rotations for the right– and left–moving electrons (these are independent because $g_{1j} = 0$), whereas the last SU(2) acts in an "orbital" space spanned by the spinors $\Psi_{Rs} = (\psi_{Rs0}, \psi_{Rs\pi})^T$ and $\Psi_{Ls} = (\psi_{Ls\pi}, \psi_{Ls0})^T$.

There is a gap m_0 in the single–particle (kink) excitation spectrum. The two–particle and particle–hole spectra can be analyzed rewriting the $\rho-$ and $\sigma-$ fields in terms of Ising model variables. [18] This then shows that excitations of orbital singlet type are massless. In the particle–particle channel this leads to powerlaw decay of d–wave pairing correlations, both in the spin singlet and triplet channels. In the particle–hole channel, the powerlaw decay occurs for CDW_π and SDW_π correlations which are out of phase between the two chains. The exponent of the powerlaw is entirely determined by the total charge degrees of freedom ($\rho+$). For $N \leq 4$ the Gross–Neveu model does not show two–kink bound states. [11] Consequently, one easily checks that all other particle–hole or two–particle excitations have the same gap $2m_0$, and the corresponding correlation functions decay exponentially with distance. These states are all triplets under the orbital SU(2).

Finally, on the second critical line $g_1 = 2g_2$ there is again a self–duality in the $\sigma-$ sector, but now the $\sigma+$ degrees of freedom also have a gap. Here $\tilde{g}^0_{11} = \tilde{g}^0_{13} = \tilde{g}^0_{21} = -\tilde{g}^0_{23} = -1/6, \tilde{g}^0_{12} = 1/3, \tilde{g}^0_{22} = 0$. We again find a Gross–Neveu model, but now with $N = 5$. The kink excitations now form a quartet, leading again to a gap m_0 in the single particle Green functions. Arranging these quartets as $\Psi_R = (\psi_{R\uparrow 0}, \psi_{R\downarrow 0}, i\psi_{R\uparrow\pi}, i\psi_{R\downarrow\pi})^T$, and $\Psi_L = (i\psi_{L\uparrow\pi}, i\psi_{L\downarrow\pi}, \psi_{L\uparrow 0}, \psi_{L\downarrow 0})^T$, these transform as SO(5) spinors, and the Hamiltonian with these coupling constants given by then is explicitly SO(5) invariant.

Contrary to the SO(3) case, there are now two–kink bound states at energy $\sqrt{3}m_0$, in addition to unbound two–kink states at $2m_0$. These bound states occur in the vector (**5**) part of the SO(5) decomposition $\mathbf{4} \otimes \mathbf{4} = \mathbf{1} \oplus \mathbf{5} \oplus \mathbf{10}$. In particular, in the particle–hole channel, the scalar (**1**) excitations are massless and occur in the OAF channel, and the bound vector excitations are the same as those in the particle–hole channel in the SU(4) case, with OAF removed. All other particle–hole excitations, occurring in the antisymmetric tensor (**10**) representation, have mass $2m_0$. In the particle–particle channel the same decomposition applies, however now the massless scalar corresponds to d–wave singlet pairing, the bound vector consists of s–wave triplet pairing and two singlet paired states with transverse momentum π, and the **10** contains all other pairing states.

The energy levels in the interesting region $0 < g_1 < 2g_2$, corresponding to repulsive interactions, and on the adjacent critical lines are shown in fig.2. In the SO(N) language, these are the discrete energy levels at $q = 0$ or $q = 2k_F$, however in our original model the presence of the massless $\rho+$ mode transforms these into lower edges of continua. Near this edge, the correlation functions have singularities of the form $(\omega - \omega_0)^{-\nu}$, where in weak coupling $\nu \approx 1/2$ and for the lowest (massless) levels $\omega_0 = 0$ and $\nu \approx 3/2$.

The full symmetries are only exact for asymptotically weak coupling. At any finite coupling, corrections to the asymptotics of eqs.(3) will lead to splittings in some multiplets. This is schematically indicated for the bound multiplets in fig.2. A more dramatic consequence of the corrections occurs in the close vicinity of the critical lines: for example, in order to obtain the massless CDW_π correlations of the SO(3) case, the correction terms have to form a bound state out of the continuum of the SO(6) spectrum and to "pull it down" all the way to masslessness as the SO(3) critical line is approached (as indicated by the broken lines in fig.2). This is in particular of interest for the case of very weakly coupled chains where renormalizations from the one–dimensional region $E > t_\perp$ lead to initial conditions for eqs.(3) obeying $g_{1i} \ll g_{2i}$. A detailed discussion is beyond the scope of the present paper.

Other parts of the phase diagram, fig.1, can be discussed along similar lines. Away from the critical lines one always finds an SO(6) symmetry. Of some interest is the SCs region, which contains the Hubbard model with attractive interaction which is numerically quite accessible and might serve as a testing ground. Here again a sixfold degenerate low–lying state exists, however now composed of the three $CDW_{0,\pi}$ states and the spin ne-



FIG. 2. Energy levels of the SO(N) models for different particle–hole and particle–particle states in the region $0 \leq g_1 \leq 2g_2$ of the phase diagram, fig.1. Horizontal lines indicate discrete energy levels, shaded areas two–particle continua. In the SO(5) case, the bound state at $\sqrt{3}m_0$ also appears in the two–particle channel, as discussed in the text. Labels indicate in which correlation function the corresponding level appears. The lowest levels in each column indicate massless correlations. TSCd: triplet d–wave superconductor; (S)CDW$_q$: (spin) charge density wave in phase ($q = 0$) and out of phase ($q = \pi$) between chains.

matic (the spin triplet analogue of the orbital antiferromagnet).

In conclusion, I have shown that in the weak–interaction limit the generic doped spin ladder model exhibits an SO(6) symmetry in its spectrum, weakly broken by corrections to scaling. The most direct consequence is that there is a sixfold degenerate *extended π–mode* in the d–wave superconducting state which contains both spin and charge excitations. This is to be compared with the SO(5) model [13] (where the symmetry is broken to SO(3) by doping) and only a triply degenerate spin π–mode is predicted. Approximately equal charge and spin excitation energies have been observed in numerical calculations [19] on t–J ladders, providing an indication that the present results may be relevant even for strongly correlated systems. To which extent the present results can be generalized to genuinely two–dimensional situations remains to be studied.

[1] M. Uehara *et al.*, J. Phys. Soc. Jpn. **65**, 2764 (1996).
[2] C. Kane and E. J. Mele, Phys. Rev. Lett. **78**, 1932 (1997), cond-mat/9608146.
[3] E. Dagotto and T. M. Rice, Science **271**, 618 (1996), cond-mat/9509181.
[4] M. Fabrizio, Phys. Rev. B **48**, 15838 (1993).
[5] H. J. Schulz, Phys. Rev. B **53**, R2959 (1996), cond-mat/9412098.
[6] L. Balents and M. P. A. Fisher, Phys. Rev. B **53**, 12133 (1996), cond-mat/9503045.
[7] E. Dagotto, J. Riera, and D. J. Scalapino, Phys. Rev. B **45**, 5744 (1992).
[8] C. A. Hayward *et al.*, Phys. Rev. Lett. **75**, 926 (1995), cond-mat/9504018.
[9] R. M. Noack, S. R. White, and D. J. Scalapino, Phys. Rev. Lett. **73**, 882 (1994), cond-mat/9409065.
[10] H.-H. Lin, L. Balents, and M. P. A. Fisher, 1998, cond-mat/9801285.
[11] A. B. Zamolodchikov and A. B. Zamolodchikov, Ann. Phys. **120**, 253 (1979).
[12] N. Andrei and J. H. Lowenstein, Phys. Lett. B **90**, 106 (1980).
[13] S.-C. Zhang, Science **275**, 1089 (1997), cond-mat/9610140.
[14] D. J. Scalapino, S.-C. Zhang, and W. Hanke, Phys. Rev. B to appear (1998), cond-mat/9711117.
[15] C. M. Varma and A. Zawadowski, Phys. Rev. B **32**, 7399 (1985).
[16] D. J. Gross and A. Neveu, Phys. Rev. D **10**, 3235 (1974).
[17] R. Shankar and E. Witten, Nucl. Phys. B **141**, 349 (1978).
[18] D. G. Shelton, A. A. Nersesyan, and A. M. Tsvelik, Phys. Rev. B **53**, 8521 (1996), cond-mat/9508042.
[19] D. Poilblanc, D. J. Scalapino, and W. Hanke, Phys. Rev. B **52**, 6796 (1995), cond-mat/9501093.